

A generalization of the Sheffer Stroke for n-valued logic.

by C.S.Scholten

For binary variables --i.e. ranging over  $\{0, 1\}$ --  $x$  and  $y$  the "Sheffer Stroke" --alias "alternative denial" or "nand"-- is the operator --or binary function-- denoted by

$$x|y$$

and (usually) defined as

$$1 - x.y$$

As  $1 - x.y = (x.y + 1) \bmod 2$

and  $x.y = \min(x, y)$

an alternative definition is:

$$x|y = (\min(x, y) + 1) \bmod 2$$

It is well-known [1] that any binary function of  $k$  binary arguments can be expressed using the Sheffer Stroke as the only primitive operator.

In the following we regard  $n$ -ary variables --i.e. ranging over  $\{0, 1, \dots, n-1\}$ -- and shall demonstrate that any  $n$ -ary function of  $k$   $n$ -ary arguments can be expressed, using as the only operator the generalization

$$x|y = (\min(x, y) + 1) \bmod n$$

In the following we shall use the notation

$$\text{suc}(x) = (x + 1) \bmod n$$

in terms of which we can rewrite:

$$x|y = \text{suc}(\min(x, y))$$

Theorem 1. The function  $\text{suc}$  --as defined above-- is expressible.

Proof.  $\text{suc}(x) = x|x$

Theorem 2. The function  $\text{pred}$  --defined by  $\text{pred}(x) = (x - 1) \bmod n$ -- is expressible.

Proof.  $\text{pred}(x) = \text{suc}^{n-1}(x)$

Theorem 3. The function  $\min$  is expressible.

Proof.  $\min(x, y) = \text{pred}(x|y)$

Note. Because  $\min(x, y, z) = \min(\min(x, y), z)$  etc.

also the minimum of more than two arguments is expressible. (End of note.)

Theorem 4. For any  $n$ -ary constant  $c$  the discriminator function  $d_c(x)$ , given by

$$\begin{aligned} d_c(x) &= n - 1 && \text{for } x = c \\ &= 0 && \text{for } x \neq c \end{aligned}$$

is expressible.

Proof. Because  $\text{suc}^{n-c}(x) = 0$  for  $x = c$   
 $> 0$  for  $x \neq c$

we have  $\min(1, \text{suc}^{n-c}(x)) = 0$  for  $x = c$   
 $= 1$  for  $x \neq c$ ,

so that  $d_c(x) = \text{pred}(\min(1, \text{suc}^{n-c}(x)))$ .

Theorem 5. For any  $n$ -ary constant  $c$  the weighted discriminator function  $w_c(x, y)$ , given by

$$\begin{aligned} w_c(x, y) &= y && \text{for } x = c \\ &= n - 1 && \text{for } x \neq c \end{aligned}$$

is expressible.

Proof.  $\min(d_c(x), \text{suc}(y)) = \text{suc}(y)$  for  $x = c$   
 $= 0$  for  $x \neq c$

and because  $\text{pred}(\text{suc}(y)) = y$ , we find

$$w_c(x, y) = \text{pred}(\min(d_c(x), \text{suc}(y)))$$

Theorem 6. The selector function, defined by

$$\text{sel}(x, y_0, y_1, \dots, y_{n-1}) = y_i \quad \text{for } x = i$$

is expressible.

Proof.  $\text{sel}(x, y_0, y_1, \dots, y_{n-1}) =$   
 $\min(w_0(x, y_0), w_1(x, y_1), \dots, w_{n-1}(x, y_{n-1}))$ .

The final induction from 2 to  $k$  arguments is straightforward.

[1] Sheffer, H.M., A set of five independent postulates for Boolean algebras, with applications to logical constants. Trans.Amer.Math.Soc. vol 14, pp.481-488